

## Tobacco Use Demographics

**Table 6. Percent (standard error) of women who reported smoking cigarettes during the 3 months before pregnancy. Ques. #28**

Maternal Characteristics	percent (s.e.)
<b>Total</b>	28.2(1.3)
<b>Race</b>	
Black	14.4(1.8)
White	36.6(1.7)
Other	25.5(8.9)
<b>Age</b>	
Less than 20	30.4(3.7)
20-29	31.1(1.7)
30+	21.6(2.2)
<b>Education</b>	
Less than High School	46.2(3.5)
High School	31.4(2.2)
More than High School	17.7(1.5)
<b>Marital Status</b>	
Married	23.7(1.5)
Other	34.0(2.1)
<b>Birthweight</b>	
LBW (<2500 grams)	30.2(3.8)
NBW (>= 2500 grams)	28.0(1.4)
<b>Medicaid Status*</b>	
Medicaid	34.6(1.8)
Other	18.0(1.7)

**Table 7. Percent (standard error) of women who reported smoking cigarettes during the last trimester of pregnancy, according to sociodemographic characteristics. Ques. #29**

Maternal Characteristics	percent (s.e.)
<b>Total</b>	17.7(1.1)
<b>Race</b>	
Black	11.6(1.6)
White	21.4(1.5)
Other	18.6(8.3)
<b>Age</b>	
Less than 20	18.5(3.1)
20-29	19.4(1.5)
30+	14.1(1.9)
<b>Education</b>	
Less than High School	33.3(3.3)
High School	19.9(1.9)
More than High School	9.1(1.2)
<b>Marital Status</b>	
Married	13.3(1.2)
Other	23.3(1.9)
<b>Birthweight</b>	
LBW (<2500 grams)	19.6(3.2)
NBW (>= 2500 grams)	17.5(1.2)
<b>Medicaid Status*</b>	
Medicaid	23.2(1.6)
Other	8.8(1.3)

**Table 8. Percent (standard error) of women who reported smoking cigarettes during the 3-6 months after delivery, according to sociodemographic characteristics. Ques #30**

Maternal Characteristics	percent (s.e.)
<b>Total</b>	23.4(1.2)
<b>Race</b>	
Black	14.2(1.8)
White	29.3(1.6)
Other	12.1(6.4)
<b>Age</b>	
Less than 20	25.9(3.5)
20-29	25.4(1.6)
30+	18.2(2.1)
<b>Education</b>	
Less than High School	39.8(3.4)
High School	26.9(2.1)
More than High School	13.3(1.4)
<b>Marital Status</b>	
Married	18.2(1.4)
Other	29.9(2.1)
<b>Birthweight</b>	
LBW (<2500 grams)	24.2(3.6)
NBW (>= 2500 grams)	23.3(1.3)
<b>Medicaid Status*</b>	
Medicaid	13.2(1.5)
Other	29.7(1.7)

\* A mother is considered "on Medicaid" if she was enrolled in Medicaid during any of the following time periods: just before pregnancy, during prenatal care or during delivery.